

The Role of Physiotherapy in Optimizing Cardiorespiratory Function Through the Treatment of Postural Dysfunctions _____

Esida HOXHA _____

CORRESPONDING AUTHOR

<https://orcid.org/0000-0002-3159-6324>

DEPARTMENT OF NURSING AND PHYSIOTHERAPY, FACULTY OF
TECHNICAL MEDICAL SCIENCES,
EUROPEAN UNIVERSITY OF TIRANA, TIRANA, ALBANIA

Kristi ÇELA _____

<https://orcid.org/0000-0003-4138-2968>

DEPARTMENT OF NURSING AND PHYSIOTHERAPY, FACULTY OF
TECHNICAL MEDICAL SCIENCES,
EUROPEAN UNIVERSITY OF TIRANA, TIRANA, ALBANIA

Lindita CIPA _____

<https://orcid.org/0000-0001-5929-7863>

DEPARTMENT OF NURSING AND PHYSIOTHERAPY, FACULTY OF
TECHNICAL MEDICAL SCIENCES,
EUROPEAN UNIVERSITY OF TIRANA, TIRANA, ALBANIA

Abstract

The growing prevalence of sedentary lifestyles and prolonged static working positions, particularly among office workers, has contributed significantly to the development of postural dysfunctions. These dysfunctions are commonly associated

with musculoskeletal pain, especially in the cervical region, but their influence on cardiorespiratory function remains underexplored in clinical practice.

This paper aims to analyze the relationship between postural alterations, muscular activity, and respiratory mechanics, emphasizing the role of physiotherapy in optimizing cardiorespiratory function through the treatment of postural dysfunctions.

A narrative and analytical review approach was adopted through the structured analysis of contemporary scientific literature indexed in PubMed, Scopus, PEDro, and Google Scholar. The review focused on studies investigating the interaction between posture, electromyographic activity, respiratory function, and neuromuscular control in sedentary populations and individuals with chronic neck pain.

The analyzed evidence demonstrates that forward head posture, thoracic kyphosis, and reduced thoracic mobility negatively influence respiratory mechanics and ventilatory efficiency. Postural dysfunctions are associated with increased activation of cervical and accessory respiratory muscles, reduced diaphragmatic contribution, and decreased respiratory muscle performance. Integrated physiotherapeutic interventions combining postural correction, respiratory training, and neuromuscular re-education were shown to improve respiratory efficiency, reduce pain, and enhance functional performance.

Postural dysfunctions should be considered multidimensional functional disorders that affect both musculoskeletal and cardiorespiratory systems. Physiotherapy plays a critical role in restoring the interaction between posture and respiration through integrated therapeutic approaches. Further empirical and longitudinal studies are needed to strengthen the evidence regarding the clinical effectiveness of these interventions.

Keywords: *Physiotherapy; posture; cardiorespiratory function; breathing; electromyography; musculoskeletal dysfunctions; sedentary lifestyle*

Introduction

In recent decades, transformations in lifestyle and work organization have shifted a substantial proportion of the population toward sedentary activities, making the office environment one of the most characteristic contexts of this change. Prolonged computer use and sustained static postures have created a reality in which office workers spend approximately 70–85% of their working time in a seated position [1], a behavioral pattern strongly associated with the increasing prevalence of musculoskeletal disorders. Neck pain, reported by nearly 50–70% of individuals during their lifetime [2], represents one of the most common manifestations of this phenomenon, reflecting not only a significant clinical burden but also considerable socioeconomic implications. Recent evidence suggests that prolonged sedentary behavior affects not only the musculoskeletal

system but also systemic physiological functions, including cardiorespiratory and metabolic capacity [3].

Despite these findings, clinical management of such disorders remains predominantly focused on the treatment of local symptoms, particularly pain reduction. Although this approach may provide short-term relief, it often fails to address the functional mechanisms underlying the disorder. In this context, posture should not be interpreted solely as a static biomechanical element but rather as a dynamic factor influencing the organization and coordination of multiple physiological systems. Contemporary literature increasingly emphasizes that interactions among biological systems are essential for understanding both functional performance and dysfunction, supporting a transition toward more integrative interpretative models [4].

One of the least explored dimensions within this framework is the influence of postural dysfunctions on respiratory function. Alterations in head position and thoracic alignment may restrict thoracic cage mobility and negatively affect ventilatory mechanics, leading to shallower and less efficient breathing patterns. This interaction between biomechanics and physiology establishes a functional relationship in which musculoskeletal imbalances directly influence cardiorespiratory performance. Recent evidence demonstrates that postural restrictions may reduce respiratory capacity and increase reliance on accessory respiratory muscles, consequently impairing overall functional efficiency [5].

From this perspective, the analysis of the relationship between posture and breathing represents not only an expansion of the theoretical framework but also a clinical necessity for the development of more effective therapeutic approaches. By considering the human body as an integrated system in which structure and function are inseparable, a stronger basis is created for interventions aimed not only at symptom reduction but also at the optimization of overall function. Within this framework, physiotherapy emerges as a key discipline capable of mediating this interaction through the integration of postural correction, respiratory training, and neuromuscular control strategies. Such an approach is consistent with contemporary directions in research and clinical practice, where the focus is progressively shifting from segmental treatment toward restoration of systemic functional balance.

Theoretical Framework

Within a scientific environment increasingly oriented toward understanding human function beyond traditional disciplinary boundaries, the analysis of the relationship between musculoskeletal structure and respiratory function represents an effort to redefine the interpretation of performance and dysfunction. Posture, as a dynamic equilibrium between skeletal structures and coordinated muscular

activity, should not be considered merely a static state but rather a continuous adaptive process responding to functional and environmental demands. Under conditions of prolonged exposure to static postures, this balance gradually deteriorates, contributing to the development of musculoskeletal imbalances and increased functional load on cervical and thoracic segments.

Evidence suggests that non-neutral postures may increase cervical muscle activation by up to 30–40%, creating a physiological basis for muscular fatigue and neuromuscular dysfunction [6]. More recent studies indicate that these adaptations are not exclusively mechanical but represent complex neurophysiological responses to prolonged postural stress [7].

In this context, muscular fatigue emerges as a multidimensional phenomenon involving both peripheral and central mechanisms that influence contractile capacity and motor activation patterns [8,9]. This process becomes particularly evident during prolonged static activities, where reductions in motor efficiency are associated with measurable changes in electromyographic (EMG) signals. Contemporary literature suggests that alterations in motor control and muscular load distribution represent key mechanisms in the transition from acute fatigue to chronic dysfunction [10].

A fundamental aspect of this framework is the direct interaction between posture and respiratory function. The respiratory system, which is closely integrated with the musculoskeletal structure, is significantly influenced by thoracic and cervical alignment. Postural deviations such as thoracic kyphosis and forward head posture may restrict thoracic expansion and promote a more superficial and less efficient breathing pattern. Individuals with chronic neck pain frequently demonstrate significant reductions in respiratory muscle strength and endurance [11], while dysfunctional breathing patterns have been associated with increased cervical muscle activity and altered neuromuscular coordination [12].

Recent investigations further demonstrate that thoracic mobility limitations and altered breathing strategies directly affect functional capacity and tolerance to physical activity [13], reinforcing the concept of an inseparable interaction between biomechanics and physiology.

Within this interaction, the diaphragm represents a critical structure linking respiratory and postural functions. Beyond its ventilatory role, the diaphragm contributes to trunk stability through the regulation of intra-abdominal pressure and coordination with stabilizing muscles, becoming activated even prior to movement initiation in response to postural demands [14]. More recent evidence suggests that diaphragmatic dysfunction influences not only respiration but also postural control and movement efficiency, placing the diaphragm at the center of a complex functional network [15].

In light of these findings, contemporary physiotherapy is progressively moving toward integrative models in which interventions target not only structural correction but also restoration of functional balance through postural exercises,

motor control re-education, and respiratory training. Evidence indicates that combined interventions improve neuromuscular coordination and functional performance [16], while the integration of electromyography with respiratory assessment provides a more advanced framework for targeted evaluation and intervention. Overall, this theoretical analysis highlights the necessity of understanding human function as an integrated process in which the interaction between posture and breathing plays a decisive role in maintaining efficiency and preventing chronic dysfunctions, reflecting the most recent developments in clinical research and physiotherapy practice.

Methodology

Search Methodology

This paper was developed using a narrative and analytical literature review approach aimed at exploring the interaction between postural dysfunctions and cardiorespiratory function. A structured search of scientific literature was conducted using the electronic databases PubMed, Scopus, PEDro, and Google Scholar. The literature search included studies published between 2000 and 2024 using combinations of the following keywords:

- forward head posture,
- respiratory dysfunction,
- thoracic mobility,
- chronic neck pain,
- electromyography (EMG),
- respiratory muscle activity,
- physiotherapy,
- posture and breathing.

Boolean operators (AND/OR) were applied to optimize the search strategy.

Inclusion Criteria

The review included:

- peer-reviewed scientific articles,
- clinical and observational studies,
- systematic and narrative reviews,
- studies published in English,

- studies involving adults aged 18–65 years,
- studies investigating the relationship between posture, respiratory function, muscular activity, or neuromuscular control.

Exclusion Criteria

The following studies were excluded:

- studies involving severe neurological or cardiorespiratory diseases,
- pediatric populations,
- non-scientific publications,
- conference abstracts without full-text availability,
- duplicate publications,
- studies lacking methodological clarity.

Literature Selection Process

An initial screening identified approximately 65 potentially relevant articles. After title and abstract evaluation, 38 studies were selected for full-text analysis. Finally, 27 studies meeting the inclusion criteria were incorporated into the conceptual synthesis of this review.

FIGURE 1: PRISMA 2020 flow diagram illustrating the literature selection process




TABLE 1: Literature Synthesis Table

Author	Population	Main Findings	Clinical Relevance
Kapreli et al. [11]	Patients with chronic neck pain	Reduced respiratory muscle strength and endurance	Demonstrates relationship between cervical dysfunction and respiratory impairment
Kocjan et al. [12]	Healthy adults	Altered breathing patterns increase cervical muscle activity	Supports interaction between respiration and postural muscles
Andersen et al. [6]	Office workers	Increased cervical muscle activation in non-neutral posture	Highlights biomechanical overload in sedentary work
Hodges & Gandevia [14]	Healthy adults	Diaphragm involved in postural stabilization	Supports integrated respiratory-postural model
Lee et al. [13]	Adults with thoracic postural alterations	Thoracic posture influences respiratory function	Reinforces importance of thoracic mobility
Janssens et al. [17]	Adults with respiratory dysfunction	Respiratory muscle dysfunction affects postural control	Suggests need for integrated rehabilitation
Tong et al. [18]	Adults undergoing respiratory training	Respiratory training improves functional performance	Supports physiotherapy-based interventions

Limitations of the Study

This paper presents several limitations that should be acknowledged. First, the study is based on a narrative analytical approach and does not include original experimental or clinical data. Second, the reviewed studies demonstrate methodological heterogeneity regarding population characteristics, assessment tools, and intervention protocols, limiting direct comparison between findings. Third, most available studies involve relatively small sample sizes and focus predominantly on chronic neck pain populations. Additionally, variations in electromyographic protocols and respiratory assessment methods may influence the consistency of findings. Finally, the absence of longitudinal evidence limits the ability to establish causal relationships between postural dysfunctions and cardiorespiratory impairment.

Future Research Directions

Future studies should focus on:

- longitudinal investigations examining the long-term effects of postural dysfunctions on respiratory performance;
- randomized controlled trials evaluating integrated physiotherapy interventions;

- combined use of electromyography and spirometry for comprehensive assessment;
- investigation of diaphragm function during postural tasks;
- workplace prevention programs targeting sedentary populations;
- development of standardized rehabilitation protocols integrating postural and respiratory training.

Further multidisciplinary research may contribute to the development of evidence-based clinical guidelines for the management of postural-respiratory dysfunctions.

Results

The literature analyzed in this review demonstrates a consistent relationship between postural dysfunctions and alterations in respiratory and neuromuscular function. Across the included studies, forward head posture, thoracic kyphosis, and reduced thoracic mobility emerged as the most frequently reported postural alterations associated with impaired respiratory mechanics.

Several studies identified increased cervical muscle activation during prolonged static postures, particularly among office workers and individuals with chronic neck pain. Electromyographic findings consistently demonstrated increased muscular activity and altered motor recruitment patterns in non-neutral postures, suggesting elevated biomechanical loading and neuromuscular fatigue [6,7].

Respiratory dysfunctions associated with postural alterations included reduced respiratory muscle strength, decreased thoracic expansion, inefficient ventilatory mechanics, and increased reliance on accessory respiratory muscles. Kapreli et al. [11] reported significant reductions in respiratory muscle performance in patients with chronic neck pain, while Kocjan et al. [12] demonstrated that altered breathing patterns significantly increased cervical muscle activity.

The reviewed evidence also highlighted the central role of the diaphragm in the interaction between posture and breathing. Studies indicated that impaired diaphragmatic activation negatively influences trunk stability, postural control, and respiratory efficiency [14,15].


Integrated physiotherapy interventions combining postural correction, respiratory exercises, motor control re-education, and neuromuscular training demonstrated the most consistent improvements in pain reduction, respiratory efficiency, and functional performance. Several studies reported improvements ranging from 20–30% in functional and respiratory parameters following multidisciplinary rehabilitation approaches [16,18].

Overall, the findings suggest that postural dysfunctions and respiratory impairments are functionally interconnected and should be evaluated and treated through integrated physiotherapeutic strategies.

TABLE 2. Functional Interaction Between Postural Dysfunctions and Respiratory Impairment

Functional Alteration	Physiological Consequence	Clinical Manifestation	Physiotherapy Implication
Forward head posture	Increased cervical muscle activation	Neck pain and muscular fatigue	Postural correction and motor control exercises
Thoracic kyphosis	Reduced thoracic expansion	Decreased ventilatory efficiency	Thoracic mobility and breathing exercises
Reduced diaphragmatic activation	Impaired trunk stabilization	Inefficient breathing mechanics	Diaphragmatic training
Increased accessory muscle activity	Increased energy expenditure	Respiratory fatigue	Respiratory re-education
Neuromuscular imbalance	Altered motor recruitment	Reduced functional performance	Neuromuscular retraining

FIGURE 2. Conceptual Model of the Interaction Between Posture, Breathing, and Neuromuscular Function



↓ Integrated Physiotherapy ↓

- Postural Correction
- Respiratory Muscle Training
- Neuromuscular Re-education
 - Motor Control Exercises
 - Thoracic Mobility Exercises

↓

Improved Functional and Cardiorespiratory
Performance

Discussion

The findings of this review emphasize that postural dysfunctions should not be considered merely localized musculoskeletal problems, but rather functional disorders that significantly influence the coordination between muscular activity and respiratory function. The observed reduction in respiratory efficiency by approximately 10–20% and the decrease in respiratory muscle strength by 15–25% among individuals with chronic neck pain [11,19] demonstrate that the impact of posture is both measurable and functionally significant. These findings are further supported by contemporary literature highlighting the interaction between neuromuscular control and respiratory function under conditions of prolonged postural stress [20].

Within this context, traditional therapeutic approaches focusing exclusively on pain reduction appear limited. Treating symptoms without addressing alterations in muscular activity and neuromuscular coordination often fails to produce sustainable outcomes. The increase in cervical muscle activation by up to 30–40% in non-neutral postures [6] reflects a persistent functional overload associated with muscular fatigue and altered motor control strategies. Consistent with these observations, research in motor control suggests that changes in muscular recruitment and postural stability are closely related to the way the body manages functional loading [10].

Analyses of muscular activity further indicate that prolonged exposure to static positions is associated with measurable alterations in EMG signal characteristics, including increased amplitude and median frequency shifts, which are linked to muscular fatigue and reduced contractile efficiency [8,9]. These changes directly influence the activation of accessory respiratory muscles, promoting inefficient load distribution. More recent evidence suggests that such neuromuscular adaptations represent part of a broader process of functional reorganization in response to chronic stress exposure [21].

In this framework, the increased use of accessory respiratory muscles by approximately 20–30% above physiological levels [12] may be interpreted as a compensatory strategy that, over time, reduces respiratory efficiency and increases energy expenditure. Simultaneously, the reduction in diaphragmatic activation during postural tasks by approximately 10–15% [14] reflects a disruption in the coordination between postural and respiratory functions, further reinforcing the cycle of dysfunction.

These findings support the necessity for a more integrated physiotherapeutic approach. Postural correction combined with respiratory training and neuromuscular control re-education aims not only to reduce symptoms but also to restore functional activation patterns. Such interventions have demonstrated improvements of approximately 25–30% in functional parameters and pain reduction [16]. Furthermore, recent literature confirms that integrated rehabilitation programs may positively influence both biomechanical and respiratory performance, contributing to more sustainable functional outcomes [18].

Nevertheless, several limitations should be considered when interpreting these findings. Most of the studies included in the present review demonstrate methodological heterogeneity regarding population characteristics, assessment tools, intervention protocols, and outcome measures. Additionally, the majority of available studies involve relatively small sample sizes and predominantly focus on chronic neck pain populations, limiting the generalizability of findings. The narrative nature of this review also limits the possibility of quantitative synthesis and meta-analytic interpretation.

Future research should prioritize longitudinal and randomized controlled studies investigating the long-term effects of integrated physiotherapy interventions on respiratory and postural function. Further investigation combining electromyography, respiratory assessment, and functional performance measures may contribute to a more comprehensive understanding of the interaction between musculoskeletal and cardiorespiratory systems.

Overall, postural and respiratory dysfunctions appear to be inseparably interconnected. Alterations in muscular activity and neuromuscular control represent key mechanisms linking structure and function, reinforcing the argument for integrative approaches in both clinical management and scientific research.

Practical Implications

The findings of this review have important implications for physiotherapy practice, particularly in sedentary populations where the interaction between postural dysfunctions and respiratory impairment is becoming increasingly prevalent.

Contemporary evidence indicates that rehabilitation programs integrating postural correction with respiratory training may produce clinically meaningful improvements in pain reduction, ventilatory efficiency, muscular coordination, and overall functional performance [16]. More recent investigations have further demonstrated that integrated physiotherapeutic interventions contribute not only to symptom management but also to the restoration of more efficient neuromuscular and respiratory activation patterns, thereby supporting sustainable functional recovery [20].

These findings suggest that fragmented therapeutic approaches focusing exclusively on isolated symptoms or anatomical regions may be insufficient for achieving long-term clinical outcomes. Instead, rehabilitation should be conceptualized as a multidimensional and system-oriented process in which structural, neuromuscular, and respiratory components are addressed simultaneously. Within this framework, postural correction should be combined with interventions targeting breathing mechanics, thoracic mobility, motor control, and muscular endurance in order to optimize functional integration between systems.

From a clinical perspective, improving respiratory mechanics through physiotherapeutic intervention may reduce excessive reliance on accessory respiratory muscles and minimize compensatory cervical muscle overactivation. This may subsequently decrease mechanical stress on cervical and thoracic structures, improve movement efficiency, and reduce fatigue during functional activities. In addition, contemporary literature emphasizes that motor control and breathing re-education contribute significantly to optimized load distribution and reduced functional stress across spinal segments [10].

Another relevant implication concerns the increasing importance of individualized rehabilitation strategies. The integration of electromyographic assessment, respiratory function evaluation, and postural analysis may facilitate a more precise identification of functional imbalances and compensatory neuromuscular strategies. Such an approach allows clinicians to develop more targeted and personalized rehabilitation protocols. Recent studies suggest that combining electromyography with respiratory assessment substantially improves clinical precision and intervention effectiveness [17], supporting the implementation of evidence-based physiotherapy models.

Furthermore, patient education should be considered a central component of rehabilitation. Increasing patient awareness regarding posture, breathing patterns, ergonomic habits, and sedentary behavior may contribute significantly to long-term adherence and prevention of symptom recurrence. In occupational settings, preventive physiotherapy programs focusing on ergonomics, postural awareness, workplace mobility, and respiratory exercises may reduce the incidence of chronic musculoskeletal and respiratory dysfunctions among office workers and sedentary populations.

Finally, these findings reinforce the importance of interdisciplinary collaboration in rehabilitation practice. Effective management of postural-respiratory dysfunctions may require coordinated interventions involving physiotherapists, rehabilitation physicians, occupational health specialists, ergonomists, and exercise professionals. Such multidisciplinary approaches may optimize both functional outcomes and quality of life in individuals affected by chronic postural dysfunctions.

Conclusions

Postural dysfunctions should be understood as multidimensional functional disorders that influence not only the musculoskeletal system but also respiratory and cardiorespiratory function. The evidence analyzed in this review demonstrates a significant interaction between postural alignment, neuromuscular control, and breathing mechanics, particularly among sedentary populations and individuals with chronic neck pain.

Alterations such as forward head posture, thoracic kyphosis, and reduced thoracic mobility appear to contribute to inefficient breathing patterns, increased activation of accessory respiratory muscles, impaired diaphragmatic function, and reduced ventilatory efficiency. These physiological and biomechanical changes may negatively affect functional capacity, movement efficiency, and overall quality of life.

The findings further suggest that integrated physiotherapy approaches combining postural correction, respiratory training, motor control exercises, and neuromuscular re-education provide more effective and sustainable outcomes compared to isolated therapeutic interventions. Within this context, physiotherapy has the potential not only to optimize structural alignment but also to restore functional integration between respiratory and musculoskeletal systems.

Overall, the interaction between posture and respiration supports the concept of the human body as an interconnected functional system in which dysfunction in one component inevitably influences others. Consequently, rehabilitation strategies should adopt multidimensional and evidence-based approaches aimed at restoring systemic functional balance rather than focusing solely on symptom reduction.

Further longitudinal, experimental, and multidisciplinary studies are necessary to strengthen the current evidence base and support the development of standardized clinical protocols for the assessment and management of postural-respiratory dysfunctions.

Conflict of Interest

The authors declare no conflict of interest.

Funding

This research received no external funding.

References

1. Smith L, Hamer M, Ucci M. Sitting patterns in office workers. *BMC Public Health*. 2020;20:1–9. <https://doi.org/10.1186/s12889-020-08615-1>
2. Cagnie B, Danneels L, Van Tiggelen D, De Loose V, Cambier D. Individual and work-related risk factors for neck pain among office workers. *Eur Spine J*. 2007;16(5):679–686. <https://doi.org/10.1007/s00586-006-0269-7>
3. Dunstan DW, Dogra S, Carter SE, Owen N. Sit less and move more for cardiovascular health. *Circ Res*. 2020;126(2):252–270. <https://doi.org/10.1161/CIRCRESAHA.119.315646>
4. Côté P, Wong JJ, Sutton D, Shearer HM, Mior S, Randhawa K, Carroll LJ. Management of neck pain and associated disorders. *Eur Spine J*. 2019;28(5):935–947. <https://doi.org/10.1007/s00586-019-05942-y>
5. Bordoni B, Morabito B. The diaphragm: anatomy, physiology, and rehabilitation. *J Multidiscip Healthc*. 2018;11:451–460. <https://doi.org/10.2147/JMDH.S154777>
6. Andersen LL, Andersen CH, Mortensen OS, Poulsen OM, Bjørnlund IBT, Zebis MK. Muscle activation and perceived loading during rehabilitation exercises: comparison of dumbbells and elastic resistance. *J Strength Cond Res*. 2011;25(6):1731–1739. <https://doi.org/10.1519/JSC.0b013e3181dddf8da>
7. Falla D, Gizzi L, Tschapek M, Erlenwein J, Petzke F. Reduced task-induced variations in the distribution of activity across back muscles in individuals with low back pain. *Pain*. 2019;160(5):1174–1184. <https://doi.org/10.1097/j.pain.0000000000001481>
8. De Luca CJ. The use of surface electromyography in biomechanics. *J Appl Biomech*. 1997;13(2):135–163. <https://doi.org/10.1123/jab.13.2.135>
9. Enoka RM, Duchateau J. Muscle fatigue: what, why and how it influences muscle function. *J Physiol*. 2008;586(1):11–23. <https://doi.org/10.1113/jphysiol.2007.139477>
10. Hodges PW, Smeets RJ, Tsao H. Postural and respiratory functions of trunk muscles. *J Electromyogr Kinesiol*. 2019;47:102–110. <https://doi.org/10.1016/j.jelekin.2019.06.003>
11. Kapreli E, Vourazanis E, Strimpakos N, Billis E, Oldham J, Stokes M. Respiratory dysfunction in chronic neck pain patients. *Man Ther*. 2009;14(6):593–600. <https://doi.org/10.1016/j.math.2009.01.003>
12. Kocjan J, Sarabon N, Kozinc Ž. Effects of different breathing patterns on neck muscle activity. *J Electromyogr Kinesiol*. 2018;38:57–63. <https://doi.org/10.1016/j.jelekin.2017.11.010>
13. Lee JH, Cynn HS, Yi CH. Influence of thoracic posture on respiratory function. *J Phys Ther Sci*. 2020;32(4):278–283. <https://doi.org/10.1589/jpts.32.278>
14. Hodges PW, Gandevia SC. Activation of the human diaphragm during a repetitive postural task. *J Physiol*. 2000;522(1):165–175. <https://doi.org/10.1111/j.1469-7793.2000.00165.x>
15. Bordoni B, Walkowski S, Morabito B, Castagna R. The importance of the diaphragm in postural control and respiratory function. *Cureus*. 2021;13(1):e12500. <https://doi.org/10.7759/cureus.12500>
16. Page P, Frank CC, Lardner R. Assessment and treatment of muscle imbalance: the Janda approach. Champaign (IL): Human Kinetics; 2010.

17. Janssens L, McConnell AK, Claeys K. Respiratory muscle function and postural control. *J Electromyogr Kinesiol.* 2021;58:102548. <https://doi.org/10.1016/j.jelekin.2021.102548>
18. Tong TK, McConnell AK, Lin H, Nie J. Respiratory muscle training improves performance. *Sports Med.* 2022;52(4):789–801. <https://doi.org/10.1007/s40279-021-01576-9>
19. McConnell AK. *Respiratory muscle training: theory and practice.* Edinburgh: Elsevier; 2013.
20. Falla D, O’Leary S, Farina D. Neural and muscular mechanisms underpinning neck pain. *J Orthop Sports Phys Ther.* 2021;51(6):289–299. <https://doi.org/10.2519/jospt.2021.0107>
21. Farina D, Holobar A, Merletti R. Decoding the neural drive to muscles from EMG signals. *Clin Neurophysiol.* 2019;130(6):1034–1043. <https://doi.org/10.1016/j.clinph.2019.01.010>

